

Image Cover Sheet

CLASSIFICATION

UNCLASSIFIED

SYSTEM NUMBER

436693



TITLE

RECORDINGS FROM SATELLITE - ALOUETTE I

System Number:

Patron Number:

Requester:

Notes:

DSIS Use only:

Deliver to: TC



JJP

225

X DRB reprint 2421

(P project D48-95-11-40)

RECORDINGS FROM SATELLITE

Alouette I

By

N. M. BRICE, R. L. SMITH, J. S. BELROSE,

and

R. E. BARRINGTON

DATE (mm)

X (Reprinted from *Nature*, Vol. 203, No. 4948, pp. 926-927, August 29, 1964)

X (Reprinted from *Nature*, Vol. 203, No. 4948, pp. 926-927,
August 29, 1964)

RECORDINGS FROM SATELLITE

Alouette I

A Very-low-frequency Plasma Resonance

AN unusual band of noise observed in the *Alouette I* satellite has been reported by Barrington and Belrose¹, who noted that the band had a sharp lower frequency cut-off which usually increased in frequency with decreasing latitude of the satellite. Since this lower cut-off frequency varies consistently with the location of the satellite, it is deduced that the observed changes in this frequency arise from spatial rather than temporal effects. Because the lower cut-off frequency is sharp and changes measurably within a few seconds, we may deduce that the horizontal field of view of the satellite for this band is at most a few tens of kilometres.

From observations of triggering of this band by both atmospherics and whistlers, Brice *et al.*² concluded that the noise was generated at the same height as the satellite. Thus the observed noise band is generated in the immediate vicinity of the satellite. Other evidence has been obtained suggesting that triggering of this band is enhanced for triggering signals propagating with large angles between the wave normal and the Earth's magnetic field. Furthermore, from examination of simultaneous very-low-frequency recordings made by the satellite and by ground-based stations, it is found that the *Alouette* hiss band is never observed on ground-based recordings.

The observation of spikes in the *Alouette* top-side sounder at the resonance frequencies for the ambient plasma³ suggests the *Alouette* hiss band arises from a similar plasma resonance at very low frequencies. The only resonance for the frequencies of interest (5-10 kc/s) at the satellite height is the lower hybrid resonance⁴ which defines a cut-off frequency for propagation transverse to the Earth's magnetic field. Other features of this resonance support the hypothesis that the lower cut-off frequency of the *Alouette* hiss band is the lower hybrid resonance for the ambient plasma.

In a subsequent paper, it will be shown that from a knowledge of the electron plasma and gyrofrequencies and the lower hybrid resonance frequency, an effective mass for the ions in the ambient plasma may be determined. Thus this hiss band may provide a powerful diagnostic tool for determining the ionic constituents of the plasma surrounding a satellite.

Assuming that the suggestion made above is correct, the observation of the *Alouette* hiss band provides, to the best of our knowledge, the first experimental confirmation of the existence of the lower hybrid resonance.

We thank Mr. John Katsufakis of Stanford for pointing out a number of the unusual features of the *Alouette* hiss band.

N. M. BRICE*
R. L. SMITH

Radioscience Laboratory,
Stanford, California.

* Present address: Faculty of Engineering, Carleton University, Ottawa, Canada.

¹ Barrington, R. E., and Belrose, J. S., *Nature*, 198, 651 (1963).

² Brice, N. M., Smith, R. L., Belrose, J. S., and Barrington, R. E. (following communication).

³ Calvert, W., and Goe, G. B., *J. Geophys. Res.*, 68 (22), 6113 (1963).

⁴ Stix, T. H., *The Theory of Plasma Waves* (McGraw-Hill Book Co., New York, 1962).

Triggered Very-low-frequency Emissions

Barrington and Belrose¹ have reported that on occasion the very-low-frequency recordings from *Alouette I* satellite (altitude, 1,000 km) showed bursts of noise triggered by atmospherics which had propagated upwards to the satellite (that is, short fractional-hop whistlers) suggesting that very-low-frequency emissions were triggered at heights of less than 1,000 km.

Since some hypotheses for the generation of very-low-frequency emissions suggested generation in or near the ionospheric *F*-region^{2,3}, while for other hypotheses generation is more likely near the top of a magnetic field line path⁴⁻⁶, it might be suggested that triggered emissions observed in *Alouette* could help to identify the mechanism responsible for generation of very-low-frequency emissions observed on ground-based recordings. However, further investigation indicates that emissions triggered by atmospherics observed in *Alouette* are not related to very-low-frequency emissions observed on ground-based recordings⁷.

Barrington and Belrose⁸ reported an unusual band of noise found in *Alouette* very-low-frequency recordings. This band showed a sharp lower frequency cut-off which increased in frequency as the satellite moved to lower latitudes. This band will be tentatively referred to as the *Alouette* hiss band. Barrington and Belrose¹ noted that the frequency of emissions triggered by atmospherics varied with the satellite location in a manner similar to that of the *Alouette* hiss band already noted here. It has now been found that these phenomena are even more closely related, in that on several occasions the emissions triggered by atmospherics were, in fact, enhancements of the *Alouette* hiss band. Furthermore, this band may also be triggered by whistlers as is illustrated in Fig. 1. The lower record of Fig. 1 is a section of the upper record (17-33 sec) shown with an expanded time-scale. Fig. 1 shows spectra obtained during a north-south pass over Stanford, California, and shows the characteristic increase of frequency with decreasing latitude and the sharp lower




Fig. 1. Very-low-frequency spectra from *Aloette I* showing triggering by atmospherics and whistlers

frequency cut-off of the hiss band. The range of L values for which the noise is seen in Fig. 1 (3.15-2.59) is consistent with that reported for the *Alouette* hiss band by Barrington *et al.*⁹. The triggering by atmospherics (propagating upwards) indicates that the hiss is generated at or below the satellite (as had been suggested by Barrington and Belrose¹) and triggering by whistlers (believed to be propagating downwards) suggests generation at or above the satellite. Thus we may deduce that the observed noise band is generated at the same height as the satellite.

Barrington *et al.*⁹ reported that examination of very-low-frequency recordings obtained during a pole-to-pole pass showed evidence that the sharp lower-frequency cut-off of the *Alouette* hiss band was very closely related to the magnetic L value and that for a given value of L the lower frequency cut-off appeared to be the same in both hemispheres. Further investigation of other recordings has shown that the close conjugate relationship with L does not always hold. A possible generation mechanism which does not require magnetic conjugacy is discussed in a companion paper by Brice and Smith⁷.

Alouette very-low-frequency recordings are made at Stanford from telemetry provided by the Canadian Defence Research Telecommunications Establishment in Ottawa.

The work at Stanford was sponsored by the National Aeronautics and Space Administration under grant *NsG* 174-61.

N. M. BRICE*
R. L. SMITH

Radioscience Laboratory,
Stanford, California.

J. S. BELROSE
R. E. BARRINGTON

✓ Defence Research Telecommunications
Establishment, (Cm)
Defence Research Board,
Ottawa.

* Present address: Faculty of Engineering, Carleton University, Ottawa, Canada.

¹ Barrington, R. E., and Belrose, J. S., Paper presented at Spring URSI Meeting, April 20-May 2, 1963 (Washington, D.C., 1963).

² Murcray, W. B., and Pope, J. H., *Phys. Rev. Letters*, 4 (1), 5 (1960).

³ Gallet, R. M., *Proc. I.R.E.*, 47 (2), 211 (1959).

⁴ Dowden, R. L., *J. Geophys. Res.*, 67 (5), 1745 (1962).

⁵ Brice, N. M. (submitted to *J. Geophys. Res.*).

⁶ Brice, N. M. (submitted to *J. Geophys. Res.*).

⁷ Brice, N. M., and Smith, R. L. (preceding communication).

⁸ Barrington, R. E., and Belrose, J. S., *Nature*, 198, 651 (1963).

⁹ Barrington, R. E., Belrose, J. S., and Keeley, D. A., *J. Geophys. Res.*, 68 (24), 6539 (1963).

JJI.

DIRECTORATE OF
SCIENTIFIC INFORMATION
SERVICES
DEFENCE RESEARCH BOARD
ROOM 4744, " " BUILDING
OTTAWA 4, ONT., CANADA

FEB - 2 1965

Date: _____
From: DSIS 1.2.65

Copy No. 1 of 7

Acc. 65/1462

#436693

- 7 - DSIS
 - Copy # 2: 1 - Ref File
 - 1 3: 1 - DATE
 - 1 4: 1 - CARDE
 - 1 5: 1 - DPhys (AE)
- (gone)